most active when recently dried, and that its efficacy was always impaired by keeping more than six months. Dr. Garden had previously made observations somewhat similar. If this be the case, we may account for its failures in the hands of those who obtain it at a distance when half a dozen years old.

Drs. Lining, Garden, and Chalmers of Carolina, are the writers who first introduced the Spigelia to notice, and who have spoken most unequivocally in its praise. Each of these physicians has represented it as an anthelmintic of superior efficacy. It appears that under certain circumstances, it is capable of operating as a cathartic, and that in these instances, the most advantage has been experienced from it. Dr. Garden says, that he had given it in hundreds of cases, and that he "never found it do much good except when it proved gently purgative." As the action of the Spigelia upon the bowels is quite uncertain, most practitioners either unite, or follow it with calomel or some purgative medicine.

We are told that the pink root, when in its most active state, if given in large quantities, induces narcotic symptoms, such as stupor, headach, dilated pupil, &c. Dr. H. Thompson, who took large doses of the root to try its effect on himself,

found that it produced an increased quickness of the pulse, drowsiness, flushing of the face and stiffness of the eyelids. Dr. Chalmers attributes to its too free use the cases of two children, who died in convulsions. Dr. Macbride informs us that its narcotic effects are seldom or never attended with danger, and that some physicians consider them an evidence of the favorable operation of the medicine. The opinion that this effect is owing to the root of some deleterious plant taken up with the Spigelia, seems to be void of foundation.

As in most other perennial plants, the root of the Spigelia possesses a greater share of activity than the herb. Of this root ten grains may be given in powder to a child four years old, twenty to one which is seven, and a drachm to an adult. If no inconvenience ensue, it may be repeated two or three times a day. If the infusion is preferred, an ounce of the root may be infused in a pint of water, and half the quantity taken by an adult or one or two spoonfuls by a child.

BOTANICAL REFERENCES.

Spigelia Marilandica, Linn. *Sp. pl.*—Curtis, *Bot. Mag. t.* 80.—Woodville, ii. *t.* 105.—Walter, *Flor. Car.* 92.—Michaux, i. 147.—Pursh, i. 139.—Elliott, i. 236.—Lonicera